

Literatur Motorbewegungsschiene

- Alfredson H, Lorenzto R (1999) Superior results with continous passive motion compared to active motion after periosteal transplantation.A retrospective study of human patella cartilage defect treatment. *Knee Surg Sports Traumatol Arthrosc* 7(4):232-8
- Aubriot JH,Guincestre JY, Grandbastien B(1993) Interet des appareils arthromoteurs dans la reeducation precoce des arthroplasties totales de genoux, *In:Revue de Chirurgie Orthopedique*,79 : 586-590
- Beaupre LA, Davies DM (2001) Exercise combined with continous passive motion or slider board therapy compared with exercise only: a randomized controlled trial of patients following total knee arthroplasty.*Phys Ther* , Apr;81(4):1029-37
- Benedikt J, Reimann M et al.(1997) Vergleichende Studie zweier Methoden der Behandlung nach Kniegelenksersatz mit bikondylärer Oberflächenersatzprothese . *Orthop. Praxis* 33:518-523
- Blauth, W (1991) Die CPM-Behandlung mit motorisierten Übungsgeräten-Historie und heutiger Stand-Praktische Hinweise.Urban und Vogel, München
- Breen, TF, Gelbermann RH, Ackerman, GN (1988) Elbow flexion contractures: treatment by anterior release and continuous passive motion. *J. Hand Surg., 13-B: 286-287*
- Breitfuß H, Muhr K et al(1991) Die Arthrolyse posttraumatischer Ellenbogensteifen. *Unfallchirurg* 94:33-39
- Chiarello CM, Gundersen L et al(1997) :The effect of continous passive motion duration and increment on range of motion in total knee arthroplasty patients. *J Orthop Sports Phys Ther* Feb;25(2):119-27
- Chylarecki C , Hierholzer G et al.(1995) Physikalische Thromboseprophylaxe mit motorisierten Sprunggelenkbewegungsschienen. *Unfallchirurgie* 21 :137-147
- Chylarecki C , Hierholzer G et al.(1997) Physikalische Thromboseprophylaxe mit einer Sprunggelenkbewegungsschiene. *Unfallchirurg* 1997
- Colwell Jr. CW, Morris BA(1992) The influence of continuous passive motion on the results of total knee arthroplasty. *Clin Orthop* 276:225-228
- Connor JC, Berk DM et al.(1995) Effects of continous passive motion following Austin bunionectomy. A pospective review.*J AM Podiatr Med Assoc* Dec;85 (12):744-8
- Coutts RD, Toth C, Kaita JH(1984) The role of continuous passive motion in the rehabilitation of the total knee patient. *In: Total Knee arthroplasty: a comprehensive approach. (Hungerford D.S., Krackow K.A., Kenna R.V., eds). Williamams & Wilkins, Baltimore: 126-132.*
- Coutts RD (1986) Continuous passive motion in the rehabilitation of the total knee patient: its role and effect. *Orthop Rev* 15:27-35
- Coutts RD, Craig EV, Mooney V et al (1988) The use of continuous passive motion in the rehabilitation of orthopaedic problems. *Contemp Orthop* 16:1-17
- Craig EV (1986) CPM in rehabilitation of surgically reconstructed shoulder; a preliminary report. Abstract presented at AAOS Meeting, New Orleans 1986

- Diehm SL (1989) The power of CPM: Healing through motion. *Continuing Care*, Vol.8, No.10
- Engström B, Sperber T, Wredmark T (1995) Continuous passive motion in rehabilitation after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol. Arthroscopy* 3:18-20
- Funk DA, Noyes FR, Grood ES, Hoffman SD (1991) Effect of flexion angle on the pressure-volume of the human knee. *Arthroscopy*; 7 (1):86-90
- Gaspar L, Farkas C et al (1997) Therapeutic value of continuous passive motion after anterior cruciate replacement. *Acta Chirurgica Hungarica*, 36 (1-4) :104-5
- Gelberman RH, Nunley JA, Osterman AL et al (1991) Influences of the protected passive mobilization interval on flexor tendon healing. A prospective randomized clinical study. *Clin Orthop* Mar; (264):189-96
- Goletz T, Henry H (1986) Continuous passive motion after total knee arthroplasty. *South. Med. J.* 79:September
- Goll SR, Lotke PA, Eckler ML (1987) Failure of continuous passive motion as prophylaxis against deep vein thrombosis after total knee arthroplasty. In: Rand J., Dorr L.D. (eds) *Total arthroplasty of the knee: proceedings of the knee society*, Aspen, Rockville M.D., p299.
- Gose JC (1987) Continuous passive motion in the post operative treatment of patients with total knee replacement: a retrospective study. *Phys Ther* 67: 39-42.
- Grassner R, Buckley MJ et al (2000) Interaction of motion and inflammation in articular cartilage effects on proteoglycan synthesis in chondrocytes In: Abstract book A 209, ITEM, 18.-20. Mai, 2000
- Hefti F, Stoll TM (1995) : Healing of ligaments and tendons; *Orthopäde* Jun; 24(3):237-45
- Jaeger T, Hassenpflug J (1991) CPM-Behandlung des Schultergelenks. *Orthopäde* 20:282-286
- James SE, Wade PJF (1987) Lateral popliteal nerve palsy as a complication of the use of a continuous passive motion knee machine – a case report. *Injury*, 18: 72-73
- Jayson MIV, Dixon ASJ (1970) Intra-articular pressure in rheumatoid arthritis of the knee. III. Pressure changes during joint use. *Ann Rheum Dis.* 29:266-8
- Johnson DP (1990) The effect of continuous passive motion on wound-healing and joint mobility after knee arthroplasty, *J. Bone Joint Surg [Am]* 72:421-426
- Johnson DP, Eastwood DM (1992) Beneficial effects of continuous passive motion after total condylar knee arthroplasty. *Ann R Coll Surg Engl* Nov; 74(6):412-6
- Jordan LR, Siegel JL, Olivo JL (1995) Early flexion routine, an alternative method of continuous passive motion. *Clin Orthop* 315:231-233
- Kaczander B (1991) The podiatric application of continuous passive motion. *J Am Podiatr Med Ass*, Vol 81, 12:631-637
- Lau SK, Chiu KY (2001) Use of continuous passive motion after total knee arthroplasty.: *J Arthroplasty* Apr; 16(3):336-9
- Lastayo PC, Wright T et al (1998) Continuous passive motion after repair of the rotator cuff. *J Bone Joint Surg* Vol.80-A, 7:1002-1011

- Levick JR(1979) The influence of hydrostatic pressure on trans-synovial movement and on capsular expansion in the rabbit knee. *J Physiol (Lond)*289:69-82
- Küsswetter W, Liebermeister R, Petzold F, Christ B (1987) Erste Erfahrungen mit einer Aktiv-Passiv-Mobilisierungsschiene für das Kniegelenk. *Aktuel Rheumatol* 12:136-140
- Kumar PJ, McPherson EJ et al (1996) Rehabilitation after total knee arthroplasty. *Clin Orthop* ;331:93-101
- Limbird T,Dennis S(1987) Synovectomy and continuous passive motion in haemophiliac patients. *J Arthroscopic Rel Surg* 3:74
- Lotke PA, Faralli VJ et al(1991)Blood loss after total knee replacement. *J Bone Joint Surg* ;73-A;No.7
- Lynch JA, Baker PL, Polly RE et al(1984) Continuous Passive Motion: A Prophylaxis for Deep Venous Thrombosis Following Total Knee Replacement, Memphis, TN, Richards Medical Co.
- Lynch AD, Bourne RB, Rorabeck CH, Rankin RN, Donald A(1988) Deep vein thrombosis and continuous passive motion after total knee arthroplasty. *J. Bone Joint Surg [Am]* 70-11
- Lynch JA, Baker PL, Polly RE et al(1990) Mechanical measures in the prophylaxis of postoperative thromboembolism in total knee arthroplasty. *Clin Orthop.* 260:24-29
- Maloney WJ, Schurmann DJ, Hangen D et al(1990) The influence of continuous passive motion on outcome in total knee arthroplasty. *Clin. Orthop* 256:162-168
- MacDonald SJ,Bourne BB,Rorabeck CH (2000) Prospective randomized clinical trial of continuous passive motion after total knee arthroplasty. *Clin Orthop Nov*;(380):30-35
- McInnes J, Larson MG, Daltroy LH et al(1992) A controlled evaluation of continuous passive motion in patients undergoing total knee arthroplasty. *JAMA* 268:1423-1428
- Merk HR, Wissel H (2002/3) Vergleichende Untersuchungen nach Schulteroperationen mit 2 postoperativen Behandlungsregimen (Vorabmitteilung an den Autor)
- O'Driscoll SW, Kumar A, Salter RB(1983) The effect of continuous passive motion on the clearance of a hemarthrosis from a synovial joint: An experimental investigation in the rabbit. *Clin. Orthop.* 176:305
- O'Driscoll SW, Salter RB(1984) The induction of neochondrogenesis in free periosteal autografts under influence of continuous passive motion: an experimental investigation in the rabbit. *J. Bone Joint Surg [Am]* 66:1248-1257
- O'Driscoll SW, Keeley FW, Salter RB(1986) The chondrogenic potential of free autogenous periosteal grafts for biological resurfacing of major full-thickness defects in joint surfaces under the influence of continuous passive motion. An experimental investigation in the rabbit. *J. Bone and Joint Surg., 68-A: 1017-1035*
- O'Driscoll SW, Keeley FW, Salter RB (1988) Durability of regenerated articular cartilage produced by free autogenous periosteal grafts in major full – thickness defects in joint surfaces under the influence of continuous passive motion. A follow-up report at one year. *J. Bone Joint Surg. (Am)* 70:595-606.
- O'Driscoll SW, Morfey BV, An KN(1990) Intra-articular pressure and capacity of the elbow. *Arthroscopy*;6:100-3.
- O'Driscoll SW (1991) Physiologische Grundlagen von “Continuous Passive Motion” (CPM). *Med Orthop. Techn.* 111: 171-174.

- Pope RO, Corcoran S, McCaul K, Howie DW(1997) Continuous passive motion after primary total knee arthroplasty. *J. Bone Joint Surg Br*; 79:914-7
- Rader CP, Kramer C et al(1998) Erfahrungen mit einer Sprunggelenksbewegungsschiene in der Thromboseprophylaxe bei Patienten nach totoalendoprothetischem Kniegelenksersatz *Z Orthop* 136:467-470
- Ritter MA, Gandolf V (1989) Continous passive motion versus physical therapy in total knee arthroplasty. *Clin Orthop*, 7:244,239-243
- Ritter Ma, Stringer EA(1979) Predictive range of motion after total knee replacement. *Clin Orthop*.143:115
- Romnes DW, Rand JA(1988) The role of continuous passive motion following total knee arthroplasty. *Clin Orthop Rel Res* 226 : 34-37.
- Rodrigo JJ, Steadman J et al (1994)Improvement of full-thickness chondral defect healing in the human knee after debridement and microfracture using continous passive motion. *Am J Knee Surg* ,Vol 7,3
- Salter, RB , Field P(1960) The effects of continuous compression on living articular cartilage. An experimental investigation. *J. Bone Joint Surg.* 42A:31
- Salter, RB, Mc Neill OR, Carbin R(1965) The pathological changes in articular cartilage associated with persistent joint deformity: An experimental investigation. *In :Gordon, D. (ed.): Studies of Rheumatoid Disease: Proceedings of the Third Canadian Conference on Research in the Rheumatic Diseases. Toronto, University of Toronto Press: 33*
- Salter RB, Simmonds DF, Malcolm BW, Rumble EJ,MacMichael D(1975) The effects of continuous passive motion on the healing of articular cartilage defects: An experimental investigation. *J. Bone Joint Surg.* 57A:570
- Salter, RB,Ogilvie-Harris DJ (1979): The healing of intraarticular fractures with continuous passive motion *In: Cooper, R. (ed.): AAOS instructional Course Lectures, St. Louis, C.V. Mosby: 102-117*
- Salter RB, Simmonds DF, Malcolm BW, Rumble EJ, MacMichael D(1980) The biological effect of continuous passive motion on the healing of full-thickness defects in articular cartilage. *J. Bone Joint Surg.* 62A:1232
- Salter RB, Bell RS(1981) The effect of continuous passive motion on the healing of partial thickness lacerations of the patellar tendon in the rabbit [abstract]. *Ann. Coll. Phys. Surg. Can.* 14:209
- Salter RB, Bell RS, Keeley, FW(1981) The protective effect of continuous passive motion on living articular cartilage in acute septic arthritis: An experimental investigation in the rabbit. *Clin. Orthop.* 159:223
- Salter RB, Minister RR(1982) The effect of continuous passive motion on a semitendinosus tenodesis in the rabbit knee [abstract]. *Orthop. Trans.* 6:292
- Salter RB, Minister RR, Clements N, Bogoch E, Bell, RS(1982) Continuous passive motion and the repair of full-thickness defects – a one-year follow-up [abstract]. *Orthop. Trans.* 6:266
- Salter RB, Hamilton HW, Wedge JH, Tile M, Torode IP, O'Driscoll SW, Murnaghan J,J Saringer JH(1983) The clinical application of basic research on continuous passive motion (CPM) for disorders and injuries of synovial joints: A preliminary report. *J. Orthop. Res.* 1:325

- Salter RB, Ogilvie-Harris DJ, Bogoch, ER(1987) Further studies in continuous passive motion [abstract]. Orthop. Trans. 2-292
- Salter RB, Rodger RM, Wilson D, Czitrom A(1987) The fate of allogenic periosteum transplanted into an osteochondral defect and subjected to continuous passive motion (CPM). An experimental investigation in the rabbit [abstract]. Clin. Med. 10(Suppl.):B127,
- Salter RB(1989) The biologic concept of continuous passive motion of synovial joints. The first 18 years of basic research and its clinical application. *Clin Orthop*, 242: 12-25
- Staubesand J, Heisterkamp T et al : Use of duplex sonograph to investigate the effect of active and passive movement on the ankle joints for promoting venous return. Clin Anat 8 , 1995 96-101
- Steadman JR, Rodkey WG et al. **Die Technik der Mikrofrakturierung zur Behandlung von kompletten Knorpeldefekten im Kniegelenk.** Orthopäde 28,1999:26-32
- Stege H , Heisterkamp Th et al.: Über die Wirkung kontinuierlicher passiver Bewegungen im oberen Sprunggelenk auf die Hämodynamik in der unteren Extremität. Phlebologie 26 (1997) 10-18
- Van Royen BJ, O'Driscoll SW, Dhert WJA, Salter RB. A comparison of the effects of immobilization and continuous passive motion on surgical wound healing in mature rabbits. Plast Reconstr Surg 1986;78: 360-6
- Ververeli P.A., Sutton D.C., Hearn S.L., Booth R.E. Jr., Hozack W.J., Rothman R.R. Continuous passive motion after total knee arthroplasty. Analysis of cost and benefits. Clin Orthop 1995 (321):208-15.
- Vince, K.G.; Kelly, M.A.; Beck, Janis; and Insall, J.N.: Continuous passive motion after total knee arthroplasty. *J. Arthroplasty*, 2:281-284, 1987.
- Walker RH, Morris BA et al.: Postoperative use of continuous passive motion, transcutaneous electrical nerve stimulation, and continuous cooling pad following total knee arthroplasty. L Arthroplasty 1991 Jun;6(2):151-6
- Wasilewski SA, Woods, LC, Torgerson Jr WR, Healy WL: Value of continuous passive motion in total knee arthroplasty. Orthopedics 13:291-295, 1990.
- Whitherow GE, Bollen SR, Pinczewski LA.: The use of continuous passive motion after arthroscopically assisted anterior cruciate ligament reconstruction: help or hindrance? Knee Surg Sports Traumatol Arthrosc 1993;1(2):68-70
- Williams J., Moran M., Thonar E., J-M.A, Salter R.B.: Continuous passive motion stimulates repair of rabbit knee articular cartilage after matrix proteoglycan loss. Clin. Orthop. 304:252 – 262, 1994
- Yashar AA, Venn- Watson E. et al. : Continuous passive motion with accelerated flexion after total knee arthroplasty. Clin Orthop. 305:38-43, 1997